

Target Space Structure of a Chiral Gauged Wess-Zumino-Witten Model

Supriya K. Kar¹ and Alok Kumar²

Institute of Physics, Bhubaneswar-751005, INDIA.

Abstract

The background for string propagation is obtained by a chiral gauging of the $SL(2, R)$ Wess-Zumino-Witten model. It is shown explicitly that the resulting background fields satisfy the field equations of the three dimensional string effective action and the target space has curvature singularity. Close connection of our solution with the three dimensional black string is demonstrated.

IP/BBSR/92-18

March '92.

¹e-mail:supriya%iopb@shakti.ernet.in

²e-mail:kumar%iopb@shakti.ernet.in

The target space structure of the gauged Wess-Zumino-Witten (WZW) model has been investigated by Witten [1] and others [2]. It was shown [1] that the vector or axial gauging of a $U(1)$ subgroup of the $SL(2, R)$ WZW model leads to the string propagation in a two dimensional target space with black hole singularity. There have been several generalizations of this result. In particular, two dimensional charged black hole [3], three dimensional black string [4] as well as four dimensional solutions with curvature singularities [5, 6] have been constructed as gauged WZW models. It has also been demonstrated that the two gaugings mentioned above are related by duality [7, 8] and they are anomaly free.

In a recent paper [9], it has been shown that there is another gauging for the two dimensional WZW model, called "chiral gauging", which is also anomaly free. Quantization of the chiral WZW model and their applications to the coset constructions have been done in ref.[9].

In this paper, we investigate the target space structure of the $SL(2, R)$ WZW model with a chiral $U(1)$ gauging. It is shown that the resulting model describes a three dimensional target space with curvature singularity. Furthermore, by an $O(2, 2)$ transformation acting on background metric ($G_{\mu\nu}$), dilaton (ϕ) and antisymmetric tensor ($B_{\mu\nu}$), it is demonstrated explicitly that our solution can be transformed to the three dimensional uncharged black string of ref.[4]. The connection to the charged black string [4] is also discussed.

We start by writing down the action for the chiral gauged Wess-Zumino-Witten ($CGWZW$) model [9] :

$$S = S^{WZW} + \frac{k}{2\pi} \int d^2z \, Tr \left[A_z^R U^{-1} \bar{\partial} U + A_{\bar{z}}^L \partial U U^{-1} + A_z^R U^{-1} A_{\bar{z}}^L U \right] \quad (1)$$

where,

$$\begin{aligned} S^{WZW} = & \frac{k}{4\pi} \int d^2z \, tr \left[U^{-1} \partial U U^{-1} \bar{\partial} U \right] \\ & + \frac{k}{12\pi} \int d^2x \, dt \, \epsilon^{ijk} Tr \left[U^{-1} \partial_i U U^{-1} \partial_j U U^{-1} \partial_k U \right] \end{aligned} \quad (2)$$

and A_z^R ($A_{\bar{z}}^L$) in eq.(1) is the $z(\bar{z})$ component of gauge field A_μ^R (A_μ^L). The classical action S is invariant under the gauge transformation [9],

$$\begin{aligned} \delta U &= v_L(\bar{z}) U - U v_R(z), \\ \delta A_{\bar{z}}^L &= -\bar{\partial} v_L(\bar{z}) \\ \text{and} \quad \delta A_z^R &= \partial v_R(z). \end{aligned} \quad (3)$$

It has been shown in ref.[9] that the theory is anomaly free. Therefore the gauge invariance is maintained at the quantum level as well. For comparison, we also write down the action for the vector $GWZW$ model [1, 10, 11] :

$$\begin{aligned} S^V = & S^{WZW} \\ & + \frac{k}{2\pi} \int d^2z \, Tr \left[A U^{-1} \bar{\partial} U + \bar{A} \partial U U^{-1} + A U^{-1} \bar{A} U + \bar{A} A \right] \end{aligned} \quad (4)$$

where in eq.(4) A and $\bar{A}$ are the two light cone components of a gauge field A_μ .

From eqs.(1) and (4), it is observed that, except for the absence of one of the terms quadratic in gauge fields, the expression for the $CGWZW$ model [9] is similar to that of the vector (or axial) one [1, 10, 11]. However it will be seen that the absence of this term makes crucial difference in the target space structure.

We now explicitly work out the case of the $SL(2, R)$ WZW model when "chiral" gauging is done with respect to the $U(1)$ subgroup generated by the currents $J_{\bar{z}}^L \equiv -\frac{i}{2} Tr (\sigma_2 U^{-1} \partial_{\bar{z}} U)$ and $J_z^R \equiv -\frac{i}{2} Tr (\sigma_2 \partial_z U U^{-1})$. Then by representing the gauge fields as, $A_{\bar{z}}^L \equiv (-\frac{i}{2} \mathcal{A}_{\bar{z}}^L \sigma_2)$, $A_z^R \equiv (-\frac{i}{2} \mathcal{A}_z^R \sigma_2)$ and the sigma model field U as,

$$U \equiv \exp \left(\frac{i}{2} \phi_L \sigma_2 \right) \exp \left(\frac{r}{2} \sigma_1 \right) \exp \left(\frac{i}{2} \phi_R \sigma_2 \right) \quad (5)$$

one obtains [10],

$$S^{WZW} = \frac{k}{4\pi} \int d^2 z \left(-\frac{1}{2} \partial \phi_L \bar{\partial} \phi_L - \frac{1}{2} \partial \phi_R \bar{\partial} \phi_R + \frac{1}{2} \partial r \bar{\partial} r - \cosh r \partial \phi_R \bar{\partial} \phi_L \right), \quad (6)$$

and

$$S = S^{WZW} + \frac{k}{4\pi} \int d^2 z \left[\mathcal{A}_z^R (\bar{\partial} \phi_R + \cosh r \bar{\partial} \phi_L) + \mathcal{A}_{\bar{z}}^L (\partial \phi_L + \cosh r \partial \phi_R) - \mathcal{A}_{\bar{z}}^L \mathcal{A}_z^R \cosh r \right]. \quad (7)$$

For comparison, we again write the $GWZW$ action, eq.(4), for the vector gauging in a similar form [10] :

$$S^V = S^{WZW} + \frac{k}{4\pi} \int d^2 z \left[\mathcal{A} (\bar{\partial} \phi_R + \cosh r \bar{\partial} \phi_L) + \bar{\mathcal{A}} (\partial \phi_L + \cosh r \partial \phi_R) - \bar{\mathcal{A}} \mathcal{A} (\cosh r + 1) \right] \quad (8)$$

It is seen that the coefficients of the last term in the RHS. of eqs.(7) and (8) are different. As pointed out earlier, the difference is due to different gauge choices in the two cases. Now, as in ref.[11], we decouple the gauge fields by a field redefinition :

$$\begin{aligned}\mathcal{A}_z^R &\equiv \hat{\mathcal{A}}_z^R + \frac{(\partial\phi_L + \cosh r \partial\phi_R)}{\cosh r} \\ \mathcal{A}_{\bar{z}}^L &\equiv \hat{\mathcal{A}}_{\bar{z}}^L + \frac{(\bar{\partial}\phi_R + \cosh r \bar{\partial}\phi_L)}{\cosh r}\end{aligned}\quad (9)$$

and obtain the action for the $CGWZW$ model as,

$$\begin{aligned}S = & \frac{k}{4\pi} \int d^2z \left(\frac{1}{2} \partial\phi_L \bar{\partial}\phi_L + \frac{1}{2} \partial\phi_R \bar{\partial}\phi_R \right. \\ & \left. + \frac{1}{2} \partial r \bar{\partial} r + \frac{1}{\cosh r} \partial\phi_L \bar{\partial}\phi_R - \hat{\mathcal{A}}_{\bar{z}}^L \hat{\mathcal{A}}_z^R \cosh r \right).\end{aligned}\quad (10)$$

By comparing this action with the following one, describing the string propagation :

$$S = \frac{1}{2\pi} \int d^2z [(G_{\mu\nu} + B_{\mu\nu}) \partial x^\mu \bar{\partial} x^\nu], \quad (11)$$

one obtains the background metric ($G_{\mu\nu}$) and antisymmetric tensor ($B_{\mu\nu}$). For example from eqs.(10) and (11), by defining an invariant distance :

$$ds^2 = \left(-\frac{k}{4} \right) \left[-dr^2 - d\phi_L^2 - d\phi_R^2 - \frac{2}{\cosh r} d\phi_L d\phi_R \right], \quad (12)$$

we get

$$G = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & -\frac{1}{\cosh r} \\ 0 & -\frac{1}{\cosh r} & -1 \end{pmatrix}, \quad (13)$$

and similarly

$$B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{\cosh r} \\ 0 & \frac{1}{\cosh r} & 0 \end{pmatrix}. \quad (14)$$

As in ref.[1], given $G_{\mu\nu}$ and $B_{\mu\nu}$, eqs.(13) and (14), the background value for the dilaton ' ϕ ' can be determined by solving the field equations of the string effective action. This procedure is followed later in the paper. The resulting dilaton background is,

$$\phi = -\ln \cosh r + \text{const.} \quad (15)$$

It can also be shown that the background dilaton of eq.(15) can be generated from eq.(10) by doing the gauge field integration in the corresponding path integral and by regularizing a $\det \left(-\frac{2\pi^2}{k \cosh r} \right)$ factor as in ref.[7].

We now show that $G_{\mu\nu}$, $B_{\mu\nu}$ and ϕ fields obtained above are in fact the solutions of the string effective action. A characteristic feature of the backgrounds in eqs.(13) - (15) is that they are dependent only on a single coordinate. String effective action with such background fields has been studied extensively by Meissner and Veneziano [12]. They worked with time dependent backgrounds, however their formalism can be applied to our case by replacing t with r in their paper. Following ref.[12], it can be shown that for the backgrounds of the type discussed in the previous paragraph, G and B can always be brought to the form,

$$G \equiv \begin{pmatrix} -1 & 0 \\ 0 & \mathcal{G} \end{pmatrix} ; \quad B \equiv \begin{pmatrix} 0 & 0 \\ 0 & \mathcal{B} \end{pmatrix} \quad (16)$$

where the first entry in a row or column of matrices G and B in eq.(16) denotes the r coordinate. To the lowest order in α' , the string tension, the complete set of field equations following from the string effective action for

this case are :

$$(\dot{\Phi})^2 - \frac{1}{4}Tr[(\mathcal{G}^{-1}\dot{\mathcal{G}})(\mathcal{G}^{-1}\dot{\mathcal{G}})] + \frac{1}{4}Tr[(\mathcal{G}^{-1}\dot{\mathcal{B}})(\mathcal{G}^{-1}\dot{\mathcal{B}})] - V = 0, \quad (17)$$

$$(\dot{\Phi})^2 - 2\ddot{\Phi} + \frac{1}{4}Tr[(\mathcal{G}^{-1}\dot{\mathcal{G}})(\mathcal{G}^{-1}\dot{\mathcal{G}})] - \frac{1}{4}Tr[(\mathcal{G}^{-1}\dot{\mathcal{B}})(\mathcal{G}^{-1}\dot{\mathcal{B}})] - V + \frac{\partial V}{\partial \Phi} = 0, \quad (18)$$

$$-\dot{\Phi}\dot{\mathcal{G}} + \ddot{\mathcal{G}} - \dot{\mathcal{G}}\mathcal{G}^{-1}\dot{\mathcal{G}} - \dot{\mathcal{B}}\mathcal{G}^{-1}\dot{\mathcal{B}} = 0, \quad (19)$$

and

$$-\dot{\Phi}\dot{\mathcal{B}} + \ddot{\mathcal{B}} - \dot{\mathcal{B}}\mathcal{G}^{-1}\dot{\mathcal{G}} - \dot{\mathcal{G}}\mathcal{G}^{-1}\dot{\mathcal{B}} = 0, \quad (20)$$

where

$$\Phi \equiv \phi - \ln \sqrt{\det \mathcal{G}} \quad (21)$$

and dots denote the derivative with respect to the r coordinate. By comparing eqs.(13) and (14) with eq.(16) we have, for our solutions,

$$\mathcal{G} = - \begin{pmatrix} 1 & \frac{1}{\cosh r} \\ \frac{1}{\cosh r} & 1 \end{pmatrix}, \quad \mathcal{B} = - \begin{pmatrix} 0 & \frac{1}{\cosh r} \\ -\frac{1}{\cosh r} & 0 \end{pmatrix}. \quad (22)$$

This implies,

$$\mathcal{G}^{-1} = - \coth^2 r \begin{pmatrix} 1 & -\frac{1}{\cosh r} \\ -\frac{1}{\cosh r} & 1 \end{pmatrix}, \quad (23)$$

$$Tr (\mathcal{G}^{-1}\dot{\mathcal{G}})^2 = 2 \left(\frac{1}{\sinh^2 r \cosh^2 r} + \frac{1}{\sinh^2 r} \right), \quad (24)$$

$$Tr (\mathcal{G}^{-1}\dot{\mathcal{B}})^2 = 2 \left(\frac{1}{\sinh^2 r \cosh^2 r} - \frac{1}{\sinh^2 r} \right) \quad (25)$$

and

$$\dot{\Phi} \equiv \dot{\phi} - \frac{1}{2} \left(\frac{\det \dot{\mathcal{G}}}{\det \mathcal{G}} \right) = - \coth r. \quad (26)$$

Using eqs.(24) - (26) it is observed that the field eq.(17) is satisfied for a constant potential $V = 1$. It can be checked that this is the same value as

for the ungauged WZW model. Also, since $\ddot{\Phi} = \frac{1}{\sinh^2 r}$, eq.(18) is satisfied for the same value $V = 1$. Eqs.(19) and (20) can be satisfied by using the following expressions :

$$-\dot{\Phi}\dot{\mathcal{G}} = \frac{1}{\cosh r} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad -\dot{\Phi}\dot{\mathcal{B}} = \frac{1}{\cosh r} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (27)$$

$$\ddot{\mathcal{G}} = \left(\frac{2}{\cosh^3 r} - \frac{1}{\cosh r} \right) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (28)$$

$$\ddot{\mathcal{B}} = \left(\frac{2}{\cosh^3 r} - \frac{1}{\cosh r} \right) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (29)$$

$$\dot{\mathcal{G}}\mathcal{G}^{-1}\dot{\mathcal{G}} = \frac{1}{\cosh^3 r} \begin{pmatrix} -\cosh r & 1 \\ 1 & -\cosh r \end{pmatrix}, \quad (30)$$

$$\dot{\mathcal{B}}\mathcal{G}^{-1}\dot{\mathcal{B}} = \frac{1}{\cosh^3 r} \begin{pmatrix} \cosh r & 1 \\ 1 & \cosh r \end{pmatrix} \quad (31)$$

and

$$\dot{\mathcal{B}}\mathcal{G}^{-1}\dot{\mathcal{G}} = \frac{1}{\cosh^3 r} \begin{pmatrix} -\cosh r & 1 \\ -1 & \cosh r \end{pmatrix} = -(\dot{\mathcal{G}}\mathcal{G}^{-1}\dot{\mathcal{B}})^T. \quad (32)$$

Therefore we have explicitly shown that the solution, eqs.(13) - (15), satisfy the field equations of the three dimensional string effective action. The scalar curvature R can be computed as [12],

$$R = 2 \left[\frac{\partial^2}{\partial r^2} (\ln \sqrt{\det \mathcal{G}}) \right] + \left[\frac{\partial}{\partial r} (\ln \sqrt{\det \mathcal{G}}) \right]^2 + \frac{1}{4} \text{Tr} (\mathcal{G}^{-1}\dot{\mathcal{G}})^2 \quad (33)$$

and we find for our case $R = -\frac{7}{2} \frac{1}{\cosh^2 r}$. Therefore the target space has curvature singularity. However it does not occur for any real value of r in the coordinate system of our choice.

Next we show that the solutions, eqs.(13) - (15), can be transformed to a three dimensional uncharged black string by an $O(2, 2)$ transformation acting on $\mathcal{G}$, $\mathcal{B}$ and Φ .

It was demonstrated in ref.[12] that the string effective action, as well as the equations of motion, (17) - (20), in space-time dimension $D (= d + 1)$ have an invariance under an $O(d, d)$ transformation acting on a matrix,

$$M \equiv \begin{pmatrix} \mathcal{G}^{-1} & -\mathcal{G}^{-1}\mathcal{B} \\ \mathcal{B}\mathcal{G}^{-1} & \mathcal{G} - \mathcal{B}\mathcal{G}^{-1}\mathcal{B} \end{pmatrix} \quad (34)$$

as $M \rightarrow \tilde{M} = \Omega M \Omega^T$ and $\Phi \rightarrow \Phi$, where Ω is an $O(d, d)$ matrix satisfying $\Omega^T \eta \Omega = \eta$ and $\eta \equiv \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$. Therefore, new solutions ($\tilde{M}$) can be generated from the old ones (M) through this transformation. To show the relation to the uncharged black string, we apply this procedure to our case. We find that for $\mathcal{G}$ and $\mathcal{B}$ as in eqs.(22) and for an $O(2, 2)$ matrix Ω :

$$\Omega \equiv \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & 0 & -\frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 & \frac{1}{\sqrt{2}} \end{pmatrix} \quad (35)$$

we have,

$$\tilde{M} = \begin{pmatrix} \tilde{\mathcal{G}}^{-1} & 0 \\ 0 & \tilde{\mathcal{G}} \end{pmatrix} \quad (36)$$

where

$$\tilde{\mathcal{G}} = -\frac{(\cosh r + 1)}{\sinh^2 r} \begin{pmatrix} \cosh r & 1 \\ 1 & \cosh r \end{pmatrix}. \quad (37)$$

By comparing eqs.(34) and (36) it is noticed that in the $O(2, 2)$ transformed coordinate system, the antisymmetric tensor $\mathcal{B}$ is absent. Since Φ

remains unchanged under this transformation, therefore we get

$$\dot{\phi} - \frac{1}{2} \left(\frac{\det \dot{\tilde{\mathcal{G}}}}{\det \tilde{\mathcal{G}}} \right) = - \coth r$$

which implies, using $(\det \tilde{\mathcal{G}}) = \coth^2 \frac{r}{2}$,

$$\tilde{\phi} = -\ln \sinh^2 \left(\frac{r}{2} \right) + \text{const.} \quad (38)$$

To show that $\tilde{\mathcal{G}}$ and $\tilde{\phi}$ in eqs.(37) and (38) are in fact the three dimensional uncharged black string, we now diagonalize $\tilde{\mathcal{G}}$ by an orthogonal transformation :

$$\tilde{\mathcal{G}}' \equiv O \tilde{\mathcal{G}} O^T = \begin{pmatrix} -\coth^2(\frac{r}{2}) & 0 \\ 0 & -1 \end{pmatrix} \quad (39)$$

where $O = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$.

From the above results we see that the three dimensional metric G of eq.(13) is transformed, under combined $O(2,2)$ (Ω) and orthogonal transformations (O), to

$$\tilde{G}' = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -\coth^2(\frac{r}{2}) & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad (40)$$

The invariant distance corresponding to the metric $\tilde{G}'$ is therefore,

$$ds^2 = \left(\frac{k}{4} \right) \left[dr^2 + \coth^2 \left(\frac{r}{2} \right) dx^2 + dy^2 \right] \quad (41)$$

where x and y are the corresponding transformed coordinates.

It is now recognised that the metric in eq.(41) represents a target space which is the cross product of the Euclidean two dimensional "dual" black hole

[1, 10] with a flat one dimensional coordinate. The solution for the dilaton in eq.(38) is also the one corresponding to the "dual" two dimensional black hole. From the three dimensional point of view, as pointed out in ref.[4] , this solution can be interpreted as the uncharged "dual" black string solution.

In fact our original solution, eqs.(13) and (14), for G and B , is also related to the charged black string even prior to the $O(2, 2)$ transformation. If we just diagonalize the G in eq.(13) we get the invariant distance :

$$ds^2 = \left(\frac{k}{4}\right) \left[dr^2 + \left(\frac{1}{\cosh r} + 1\right) dx^2 + \left(\frac{1}{\cosh r} - 1\right) dy^2 \right] \quad (42)$$

and the antisymmetric tensor :

$$B = \frac{1}{\cosh r} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (43)$$

Then by defining $\rho = \cosh r$ and $y = i t$ we have,

$$ds^2 = \left(\frac{k}{4}\right) \left[- \left(1 - \frac{1}{\rho}\right) dt^2 + \left(1 + \frac{1}{\rho}\right) dx^2 + \left(1 - \frac{1}{\rho}\right)^{-1} \left(1 + \frac{1}{\rho}\right)^{-1} \frac{d\rho^2}{\rho^2} \right], \quad (44)$$

$$B = \frac{i}{\rho} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} \quad (45)$$

and

$$\phi = - \ln \rho + \text{const.} \quad (46)$$

This solutions matches with the one in ref.[4] for the charged black string if one takes $\lambda = -\frac{1}{2}$ and defines $\rho = 2 \hat{r}$ in eqs.(12) and (13) of this reference. However the B field in this case is imaginary. This field can be made real by

keeping t space-like. The value $\lambda = -\frac{1}{2}$ in ref.[4] implies that the free boson 'x' of the $[SL(2, R) \times R]$ WZW model is time like.

To conclude, in this paper the background graviton, dilaton and antisymmetric tensor fields resulting from the chiral $U(1)$ gauging of the $SL(2, R)$ WZW model were obtained. It was shown that the backgrounds satisfy the field equations of the string effective action and target space has curvature singularity. Relation of our solution with the three dimensional black string has been shown. An interesting aspect of the chiral gauging is that the resulting model has the same target space dimension as the original one. Whereas due to the vector or axial gauging the target space dimension is reduced. It will be interesting to examine the duality properties of the $CGWZW$ models as in the vector and axial cases. Also, since the chiral gauging is left-right asymmetric in nature, it should be possible to obtain exact conformal field theory solutions for the heterotic strings in nontrivial background using $CGWZW$ models.

References

- [1] E. Witten, Phys. Rev. **D44**, 314 (1991)
- [2] I. Bars, Univ. of Southern California, Los Angeles preprint USC-91-HEP-B3 (1991); E. Martinec and S. Shatashvili, EFI preprint EFI-91-22 (1991); M. Bershadsky and D. Kutasov PUPT-1261, HUTP-91/A 024 (1991).
- [3] N. Ishibashi, M. Li and A. R. Steif, Phys. Rev. Lett. **67**, 3336 (1991); S. Khastgir and A. Kumar, Mod. Phys. Lett. **A6**, 3365 (1991).
- [4] J. H. Horne and G. T. Horowitz, Univ. of California, Santa Barbara preprint UCSBTH-91-39, July 1991.
- [5] I. Bars and K. Sfetsos, Univ. of Southern California, Los Angeles preprint USC-91-HEP-B6 (1991).
- [6] P. Horava, Enrico Fermi Institute preprint EFI-91-57 (1991).
- [7] E. B. Kiritsis, Mod. Phys. Lett. **A6**, 2871 (1991).
- [8] M. Rocek and E. Verlinde, Institute for Advanced Study preprint IASSNS-HEP-91/68 (1991).
- [9] S. Chung and S. H. Henry Tye, Newman Laboratory preprint CLNS-91-1127.

- [10] R. Dijkgraaf, E. Verlinde and H. Verlinde, Institute for Advanced Study preprint IASSNS-HEP-91/22 (1991).
- [11] S. Nojiri, Phys. Lett. **B274**, 41 (1992).
- [12] K. A. Meissner and G. Veneziano, Phys. Lett. **B267**, 33 (1991); Mod. Phys. Lett. **A6**, 3397 (1991).